

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number
WO 03/080566 A2

(51) International Patent Classification⁷: C07C 259/06,
C07D 207/46, C07C 59/347, C07K 5/06, C07C 309/17,
C07F 5/02, A61K 31/194, 31/16, 31/33

Peter, John [GB/GB]; Wellcome Trust Centre for Human
Genetics, Roosevelt Drive, Oxford OX3 7BN (GB).

(21) International Application Number: PCT/GB03/01239

(74) Agents: ELLIS-JONES, Patrick, George, Armine et al.;
J.A. Kemp & Co., 14 South Square, Gray's Inn, London
WC1R 5JJ (GB).

(22) International Filing Date: 21 March 2003 (21.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0206711.4 21 March 2002 (21.03.2002) GB

(81) Designated States (*national*): AB, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): ISIS IN-
NOVATION LIMITED [GB/GB]; Ewert House, Ewert
Place, Summertown, Oxford OX2 7SG (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and






(75) Inventors/Applicants (*for US only*): SCHOFIELD,
Christopher, Joseph [GB/GB]; Dyson Perrins Labora-
tory, University of Oxford, South Parks Road, Oxford
OX1 3QY (GB). MAXWELL, Patrick, Henry [GB/GB];
Imperial College of Science, Technology & Medicine,
Hammersmith Campus, Du Cane Road, London W12 0NN
(GB). PUGH, Christopher, William [GB/GB]; University
of Oxford, Wellcome Trust Centre for Human Genetics,
Roosevelt Drive, Oxford OX3 7BN (GB). RATCLIFFE,

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HIF HYDROXYLASE INHIBITORS



(57) Abstract: The invention provides a compound of one of the formulae (A), (B), (C), (D), (E), (F) as herein defined, or a salt thereof, for use in the treatment of a condition associated with increased or decreased HIF levels or activity, or a condition in which an increase or decrease in HIF levels or activity may be beneficial.

BEST AVAILABLE COPY

WO 03/080566 A2

HIF HYDROXYLASE INHIBITORS

Field of Invention

The present invention relates to compounds which modulate 2OG (2-oxoglutarate) dependent oxygenases, in particular prolyl hydroxylases. These may be useful as modulators of HIF (hypoxia inducible factor) alpha (HIF- α) prolyl hydroxylase.

Background of Invention

The transcription factor HIF (hypoxia inducible factor) system is a key regulator of responses to hypoxia, occupying a central position in oxygen homeostasis in a wide range of organisms. A large number of transcriptional targets have been identified, with critical roles in angiogenesis, erythropoiesis, energy metabolism, inflammation, vasomotor function, and apoptotic/proliferative responses. The system is essential for normal development, and plays a key role in pathophysiological responses to ischaemia/hypoxia. HIF is also important in cancer, in which it is commonly upregulated, and has major effects on tumour growth and angiogenesis. The HIF DNA binding complex consists of a heterodimer of α and β subunits. Regulation by oxygen occurs through hydroxylation of the α -subunits, which are rapidly destroyed by the proteasome in oxygenated cells. This involves binding of HIF- α -subunits by the von Hippel-Lindau tumour suppressor protein (pVHL), with pVHL acting as the, or part of the, recognition component for a ubiquitin ligase that promotes ubiquitin dependent proteolysis through interaction with a specific sequence or sequences in HIF- α -subunits. In hypoxia, this process is suppressed, so stabilizing HIF- α and permitting transcriptional activation via the HIF α , β .

Disclosure of the Invention

In our British Application No. 0118952.1 we disclose a polypeptide comprising:

- (a) the amino acid sequence of SEQ ID NO: 2, 4 or 6 having HIF hydroxylase activity;
- (b) a variant thereof having at least 60% identity to the amino acid

sequence of SEQ ID NO: 2, 4 or 6 and having hydroxylase activity; or

(c) a fragment of either thereof having HIF- α hydroxylase activity.

Preferably, the polypeptides have prolyl hydroxylase activity and require Fe(II) for activity.

5 They are related by sequence to non-haem oxygenases for which crystal structures are known, e.g. proline-3-hydroxylase (Clifton et al, Eur. J. Biochem, 2001, 268, 6625-6636).

It also discloses polynucleotides which encode the polypeptides as well as expression vectors comprising the polynucleotide and antibodies capable of specifically binding the polypeptide. We also disclose assays for identifying
10 modulators of the activity of the HIF hydroxylase as well as the use of modulators such as inhibitors of the activity of the peptides in the treatment of a condition or disease associated with altered HIF levels with respect to healthy (or normal) levels, and the treatment of conditions where an alteration in the HIF levels or activity
15 would be beneficial.


Inhibitors of the 2-OG dependent enzyme collagen prolyl-4-hydroxylase (CPH) are well known in the art and have been previously proposed for use in the treatment of lung fibrosis, skin fibrosis (scleroderma), atherosclerosis and other conditions associated with collagen biosynthesis. Inhibitors of para-
20 hydroxyphenylpyruvate oxygenase (a non-haem oxygenase employing ferrous iron as a co-factor) such as triketones are used as herbicides (Lee D. et al (1998) Pestic. Sci. 54(4) 377-384). We have disclosed that certain of these CPH inhibitors (and other components) also inhibit the biological (i.e. HPH) activity of an PHD polypeptide. A CPH inhibitor or modified CPH inhibitor which inhibits the biological activity of an
25 PHD polypeptide may be used in the treatment of a condition associated with reduced or suboptimal HIF levels or activity, or a condition in which an increase in HIF levels or activity may be beneficial, for example ischaemia, wound healing, auto-, allo-, and xeno- transplantation, systemic high blood pressure, cancer, inflammatory disorders, and metabolic disorders.

30 Various methods and uses of modulators which inhibit, potentiate, increase or stimulate hydroxylation of HIF- α by an PHD polypeptide are disclosed. The purpose

of disruption, interference with or modulation of the hydroxylation of HIF-1 α by a PHD polypeptide may be to modulate cellular functions such as angiogenesis, erythropoiesis, energy metabolism, inflammation, matrix metabolism, vasomotor function, and apoptotic/proliferative responses and pathophysiological responses to ischaemia/hypoxia, all of which are mediated by HIF α as discussed above.


Compounds which modulate 2OG oxygenases, in particular CPH may be useful as modulators of HIF prolyl hydroxylase, or may be used as 'lead' compounds which may be modified and/or optimised to develop modulators of HIF prolyl hydroxylase, in particular selective modulators are described.

Some of these compounds generally possess the formula:




where the group R^1 is capable of forming an electrostatic interaction with the sidechain of the arginine residue which, together with other residues, binds the 5-carboxylate of 2-oxoglutarate during catalysis, $A*B$ is a chain of two atoms which are, independently, carbon, oxygen, nitrogen or sulphur, which chain can be functionalised, y is 0 or 1 and $C*D$ is a chain of two atoms which are, independently, carbon, oxygen, nitrogen, or sulphur, which chain can be functionalised, A , B , C and D being linked to one another by a single and/or double and/or triple bond such that when y is 0 or 1 at least one of the atoms of which is capable of chelating with a metal group and when y is 1 said chain is attached to R^2 which is capable of chelating with a metal group. Generally at least one of A , B , C and D is not carbon. Typical chains include $C-N-C-C$ and $C-O-C-C$ and $C-C-C=O$. The chain atoms can form part of a ring.

New classes of modulators of HIF prolyl hydroxylase have been found, according to the present invention. These possess the following formulae (A) to (F)




where each of R^1 and R^5 is independently H, OH, SH, a branched or straight C_1 to C_6


alkyl chain optionally containing 1 or more eg. 2 N, S, O or P chain atoms, especially methyl, which can be functionalised, any amino acid side chain, such as alanine, phenylalanine, valine and glutamic acid, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring, such as aryloxy alkyl, A¹ is CH₂ or O, and each of R² and R³ is independently be H, OH, a branched or straight C₁ to C₆ alkyl chain optionally containing 1 or more eg. 2 N, S, O or P chain atoms which can be functionalised, optionally with 1, 2, 3, 4 or 5 halo substitutions, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring,




wherein R² is as defined above, --- is a single bond and T is CH₂ or C=O, or --- is a double bond and T is CH; A² is H or -XCO₂R⁴; X is a single bond or a branched or straight C₁ to C₆ alkyl chain, optionally containing 1 or more eg. 2 N, S, O or P chain atoms and optionally substituted by eg. halo, OH, NHR² or NHCOR⁴ where R² and R⁴ are as defined above and R⁴ represents H, a branched or straight chain C₁ to C₆ alkyl group optionally containing 1 or more eg. 2 N, S, O or P chain atoms, a 4 to 7 membered heterocyclic ring, optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, which may be fused to another ring, or a salt thereof,



where each X which may be the same or different is NH, NRⁿ, where Rⁿ is OH, a branched or straight C₁ to C₆ alkyl chain optionally containing 1 or more eg. 2 N, S, O or P chain atoms which can be functionalised, or O i.e. XR¹ is typically OH or O-alkyl having a branched or straight C₁ to C₆ alkyl chain, especially MeO,
 5 each Y, which may be the same or different, is O or S and each R¹, which may be the same or different, is as defined above,



10 where m is 0 or 1, Q represents (R¹R⁶)_xZ where x is 0, 1 or 2, R¹ is as defined above and R⁶ is as defined for R¹, and Z is P=O(OH)₂, B(OH)₂ or SO₃H, or a salt thereof, typically a sodium salt, or




where each R¹, which are the same or different, is as defined above; R¹¹ represents OH or R¹⁰ NH where R¹⁰ is HO, R¹CO or HOOC(X)_x wherein R¹ is as defined above,
 20 x is 0 or 1 and X is R¹R¹C wherein each R¹, which are the same or different, is as defined above; or R¹⁰ is an amino acid residue H₂N (R¹R¹C) CO - wherein each R¹, which are the same or different, is as defined above; n is 1 or 2 and R¹² is H or straight or branched C₁ to C₆ alkyl; or a salt thereof. Typically X is CH₂ or CHOH.

Another aspect of the invention concerns analogues of 2-oxoglutarate that act
 25 as improved (relative to 2-oxoglutarate) co-substrates for the HIF hydroxylases. Such a compound is 3-fluoro 2-oxoglutarate. Assays in which this compound replaces 2-oxoglutarate demonstrate a higher level of HIF hydroxylation than observed when using 2-oxoglutarate under analogous conditions.

These analogues possess the formula:

6



5 wherein each of Z^1 and Z^2 is independently hydrogen, SH or an electron withdrawing group such as halogen, preferably fluorine, or alkoxy such as methoxy, and R^{12} is as defined above, or a salt thereof. Preferably one of Z^1 and Z^2 is hydrogen and the other is fluorine (3-F-2-OG).

 Accordingly the present invention provides a compound of formula (A) to
 10 (F) for use in the treatment of a condition associated with increased or decreased HIF levels or activity, or a condition in which an increase or decrease in HIF levels or activity may be beneficial, as well as the use of a compound of formula (A) to (F) in the manufacture of a medicament for the treatment of such a condition.

 The said alkyl groups and chains are typically functionalised by alcohol,
 15 fluorine, thiol, a carboxylic acid, phosphonic or phosphinic acid, sulphonic acid or other chelating group, in the case of the chains typically via an alkyl group.

 In the formulae described herein, a branched or straight C_1 to C_6 alkyl chain may be a methyl, ethyl, propyl, butyl, iso-butyl, *tert*-butyl, pentyl, neopentyl, *tert*-pentyl or a primary, secondary or tertiary hexyl group. Hetero atoms such as O, S, N
 20 and P may replace one or more of the carbon atoms. Preferably the alkyl groups are methyl, the preferred heterocyclic rings are pyrrolidine and tetrahydropyran and the preferred aromatic rings are benzene, naphthalene and pyridine.

 Typically, each of R^1 and R^5 is independently H, OH, a branched or straight C_1 to C_6 alkyl chain optionally containing 1 or more N, S, O or P chain atoms, which
 25 can be functionalised, any amino acid side chain, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring.

 Typically, A^1 is CH_2 .

30 Typically, A^2 is $-\text{XCO}_2\text{R}^4$.

 Typically, R^{11} represents $R^{10}\text{NH}$ where R^{10} is $R^1\text{CO}$ or $\text{HOOC}(\text{X})_x$ wherein

R¹ is as defined above, x is 0 or 1 and X is R¹R¹C wherein each R¹, which are the same or different, is as defined above; or R¹⁰ is an amino acid residue H₂N (R¹R¹C) CO- wherein each R¹, which are the same or different, is as defined above.

Typically, each of Z¹ and Z² is independently hydrogen or an electron
5 withdrawing group.

Typically, in the compounds of formula (F), R¹² is H. Alternatively, R¹² may be straight or branched C₁ to C₆ alkyl.

The compounds of formula (A) are hydroxamates. Preferred compounds include those where R⁵ is aryloxyalkyl, especially oxyloxymethyl such as
10 phenyloxymethyl or phenylalkyloxymethyl, especially benzyloxymethyl or substituted benzyloxymethyl such as p-hydroxy benzyloxymethyl and/or where R² and/or R³ is HOCH₂.

Typical compounds include N-phenoxy-acetyl-(L)-alanine-hydroxamide (Is41) and the corresponding (D) isomer (Is43) as well as the corresponding tyrosine
15 derivatives (Is44 and 45) and L- and D-phenylglycine derivatives (Is46 and 47), along with benzo hydroxamic acid and N-phenoxyacetyl-D-phenylalanine hydroxamic acid (Is42).

These compounds can generally be prepared following the method of Walter et al., Tetrahedron 1997, 53, 7275-7290 and Biorg. Chem 1999, 27, 35-40.

20 The compounds of formula (B) are cyclic hydroxamates. Preferred compounds are those where X is a single bond or methyl and/or R² is H or phenylalkyl, especially benzyl and/or R⁴ is H or methyl. Typical compounds include (1-hydroxy-2, 5-dioxo-pyrrolidin-3-yl) acetic acid (Is52), (1-hydroxy-2, 5-dioxo-pyrrolidin-3-yl) carboxylic acid (ANU 2) and its N-benzoyloxy derivative (ANU 1)
25 along with (1-benzyloxy-2, 5-dioxo-pyrrolidin-3-yl) acetic acid (Is50) and the corresponding methyl ester (Is64), and N-hydroxy succinimide (C1). Note that Is52 (R²=H, T=C=O, X=CH₂, R⁴=H) is highly active reflecting its structural analogy with 2-oxoglutarate. These compounds can be prepared using the general procedure of Schlicht et al. (US 4446038).

30 The compounds of formula (C) are analogues of 2-oxoglutarate or oxalyl derivatives of hydroxyacetate and mercapto acetic acid. Preferred compounds

include those where X is O and/or R¹ is H or methyl. Typical compounds include dimethyl oxalylglycolate (Is10) as well as its free acid (Is14) and dimethyl oxalylthioglycolate (Is11). These compounds can be prepared following Franklin et al., J. Med. Chem 1992, 35, 2652-2658 or Kwon et al., J. Am. Chem. Soc. 1989, 111, 1854-1860.

The compounds of formula (D) are carboxylic acids which possess a phosphonic, sulphonic or boronic acid group as well as salts of these. Typically R¹ and R⁶ are hydrogen. Preferred compounds include the phosphoric acids where x is 0, 1 or 2 (C3, 4 and 5, respectively) as well as disodium 3-sulpho-propionate (Is63) and its free acid, and 3-borono-propionic acid (Is62).

The compounds of formula (E) are N-acylated amino acids or polycarboxylic acids. Typical compounds are those where R¹ is H, and/or R¹² is H or ethyl. When R¹¹ represents R¹⁰NH the compounds are typically dipeptides such that R¹⁰ is an acyl group of a natural amino acid such as glycine. Typical preferred such compounds include Asp-Gly (C18), cyclo (Asp-Gly) (C19), beta-Asp-Gly (C20), Glu-Gly (C21) and Z-Glu-Gly (C22). Other typical compounds include those where R¹⁰ is acetyl or benzoyl such as the N-acetylated derivatives of L-aspartic acid (C6) and of L-glutamic acid (C7) i.e. R¹⁰ is acetyl and N-benzoylated derivatives of glutamic acid (C15 and Is90) i.e. R¹⁰ is benzoyl. Other typical compounds include those where R¹¹ is -NHOH such as diethyl 2-(hydroxylamino)-glutarate (Is51 being the racemic form of this compound) and those where R¹¹ is OH such as 2-hydroxyglutaric acid (Is57). When R¹¹ is HOOC(X)x, X is especially CH² or CHOH. The compounds are typically citric acid (C12), tricarballic acid (C13) and succinic acid as well as the tri-methyl ester of ethane tricarboxylic acid (Is72).

The compounds of formula (F) are analogues of 2-oxoglutarate. Preferred compounds include 3-fluoro-2-oxoglutarate compounds (i.e. Z¹ is H and Z² is F) such as 3-fluoro-2-oxoglutaric acid (Is18) and the corresponding dimethyl ester (Is19).

The compounds which are acids can be present in the form of salts, such as sodium salts.

For therapeutic treatment, the compound may be used in combination with any other active substance, e.g. for anti-tumour therapy another anti-tumour

compound or therapy, such as radiotherapy or chemotherapy.

Generally, the modulator is provided in an isolated and/or purified form, i.e. substantially pure. This may include being in a composition where it represents at least about 90% active ingredient, more preferably at least about 95%, more preferably at least about 98%. Any such composition may, however, include inert carrier materials or other pharmaceutically and physiologically acceptable excipients, such as those required for correct delivery, release and/or stabilisation of the active agent. As noted below, a composition according to the present invention may include in addition to a modulator compound as disclosed, one or more other molecules of therapeutic use, such as an anti-tumour agent.

In general they take the form of compositions wherein the compound is in a mixture with a pharmaceutically acceptable carrier or diluent. The carrier may be liquid, e.g. saline, ethanol, glycerol and mixtures thereof, or solid, e.g. in the form of a tablet, or in a semi-solid form such as a gel formulated as a depot formulation or in a transdermally administerable vehicle, such as a transdermal patch. The modulator compound or composition comprising it may be formulated as the coating of a coated stent.

The invention further provides a method of treatment which includes administering to a patient compound as defined above. Exemplary purposes of such treatment are discussed elsewhere herein.

The therapeutic/prophylactic purpose of such a method or use may be the modulation of the level of HIF α in a cell by modulation, e.g. disruption or interference, of the hydroxylation of HIF α , which may occur for example at proline 402, 564 or other proline residue. Hydroxylation of HIF α promotes pVHL binding which leads to ubiquitin dependent proteolysis of HIF α as described above.

The therapeutic/prophylactic purpose may be related to the treatment of a condition associated with reduced or suboptimal or increased HIF levels or activity, or conditions where an alteration in HIF levels or activity may be beneficial such as: (i) ischaemic conditions, for example organ ischaemia, including coronary, cerebrovascular and peripheral vascular insufficiency. The therapy may be applied in two ways; following declared tissue damage, e.g. myocardial infarction (in order to

limit tissue damage), or prophylactically to prevent or ameliorate ischaemia, e.g. promotion of coronary collaterals in the treatment of angina.

- (ii) wound healing and organ regeneration.
- (iii) auto-, allo-, and xeno- transplantation.
- 5 (iv) systemic blood pressure.
- (v) cancer; HIF α is commonly up-regulated in tumour cells and has major effects on tumour growth and angiogenesis.
- (vi) inflammatory disorders.
- (vii) pulmonary arterial blood pressure, neurodegenerative disease.
- 10 (viii) metabolic disorders, e.g. diabetes.

Modulating HIF prolyl hydroxylase activity in a person, an organ, or a group of cells may be exploited in different ways to obtain a therapeutic benefit:

- (a) Non cell autonomous: The HIF system is used by cells to influence the production of substances which signal to other cells. These signals may then have effects at (i) a distant site (for example erythropoietin acts on the bone marrow) or
15 (ii) locally (angiogenic growth factors increase the local formation of blood vessels). Manipulating non cell autonomous behaviour via altering hydroxylase activity is therefore useful in the treatment of anaemia, and local ischaemia, for example in the eye, brain, heart and limbs. Many other signals that are involved in aspects of
20 physiological homeostasis may be, or are known to be, adjusted by HIF activation. Consequently altering HIF prolyl hydroxylase activity may be used to potentiate or initiate a helpful response for a therapeutic benefit, or to prevent or ameliorate a harmful response. For example, this approach can be used to alter appetite, or blood pressure in the systemic or pulmonary beds.
- 25 (b) Cell autonomous: the HIF system is also used by cells to regulate cellular metabolism, and decisions concerning differentiation, proliferation and apoptosis. Therefore manipulating the HIF system can be used to alter the viability and behaviour of cells. An increase in cell viability can be achieved by increasing HIF activation, for example in an ischaemic tissue. This approach can also be used
30 in improving pancreatic beta cell viability as a way of ameliorating diabetes, or of improving the viability or function of a group or groups of neurons in Parkinson's

disease, motorneurone disease or forms of dementia. In a different approach, the HIF signal can be manipulated to prevent a group of cells proliferating, or to promote its death or differentiation. For example transient activation of the HIF system in a malignant tumour can be used to provoke death of a substantial number of tumour cells.

Pharmaceutical Compositions

In various further aspects, the present invention thus provides a pharmaceutical composition, medicament, drug or other composition for such a purpose, the composition comprising one or more compounds of formulae (A) to (F), or derivatives thereof, the use of such an composition in a method of medical treatment, a method comprising administration of such a composition to a patient, e.g. for treatment (which may include preventative treatment) of a medical condition as described above, use of such an agent compound or substance in the manufacture of a composition, medicament or drug for administration for any such purpose, e.g. for treatment of a condition as described herein, and a method of making a pharmaceutical composition comprising admixing such an agent, compound or substance with a pharmaceutically acceptable excipient, vehicle or carrier, and optionally other ingredients.

The agent may be used as sole active agent or in combination with one another or with any other active substance, e.g. for anti-tumour therapy another anti-tumour compound or therapy, such as radiotherapy or chemotherapy.

Whatever the agent used in a method of medical treatment of the present invention, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors.

An agent or composition may be administered alone or in combination with

other treatments, either simultaneously or sequentially dependent upon the condition to be treated, e.g. as described above.

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may include, in addition to active
5 ingredient, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous or intravenous. The
10 compositions will typically be sterile.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid
15 carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability.
20 Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

Liposomes, particularly cationic liposomes, may be used in carrier
25 formulations. Examples of techniques and protocols mentioned above can be found in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed), 1980.

The substance or composition may be administered in a localised manner to a particular site or may be delivered in a manner in which it targets particular cells or tissues, for example using intra-arterial stent based delivery.

30 Targeting therapies may be used to deliver the active substance more specifically to certain types of cell, by the use of targeting systems such as antibody

or cell specific ligands. Targeting may be desirable for a variety of reasons, for example if the agent is unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.


5 The following Examples further illustrate the present invention.

Example 1

 In vitro screening of potential inhibitors of HIF modification was performed using a capture assay. A Gal/HIF-1 α /VP16 fusion protein expressing HIF-1 α residues 549-582 was prepared by IVTT (see British Application No. 0118952.1) and
10 used as a substrate in the assay. The unlabelled substrate was immunopurified on beads, washed, and aliquots incubated in the presence of RCC4 cell extract, with 100 μ M FeCl₂ and 2mM of the potential inhibitor. The inhibitors were either dissolved in DMSO or Tris as indicated. Controls, where no inhibitor but the equivalent amount of DMSO or Tris was added, were also performed. After washing,
15 the beads were assayed for their ability to interact with 35-S labelled pVHL IVTT. Hydroxylation of the fusion protein by HIF hydroxylase present in the cell extract leads to the ability to capture the labelled pVHL and the amount of labelled protein bound to the fusion protein can then be measured to determine relative HIF hydroxylase activity. Figures 1 to 5 show HIF hydroxylase activity in the presence of
20 a particular inhibitor relative to that seen in the absence of the inhibitor (the DMSO/Tris control).


CLAIMS

1. A compound of one of the formulae




- where each of R^1 and R^5 is independently H, OH, SH, a branched or straight C_1 to C_6 alkyl chain optionally containing 1 or more N, S, O or P chain atoms, which can be functionalised, any amino acid side chain, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring, A^1 is CH_2 or O, and each of R^2 and R^3 is independently H, OH, a branched or straight C_1 to C_6 alkyl chain optionally containing 1 or more N, S, O or P chain atoms which can be functionalised, optionally with 1, 2, 3, 4 or 5 halo substitutions, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring,

20



- 25 wherein R^2 is as defined above, --- is a single bond and T is CH_2 or $C=O$, or --- is a double bond and T is CH; A^2 is H or $-XCO_2R^4$; X is a single bond or a branched or straight C_1 to C_6 alkyl chain optionally containing 1 or more N, S, O or P chain atoms, and optionally substituted by halo, OH, OR^4 , NRR^2 or $NHCOR^4$ where R^2 , R^3 and R^4 are as defined above and R^4 represents H, a branched or straight chain C_1 to C_6 alkyl group, a 4 to 7 membered heterocyclic ring, optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N,
- 30

O or S atoms, which may be fused to another ring,



5


where each X which may be the same or different is NH, NRⁿ, where Rⁿ is OH, a branched or straight C₁ to C₆ alkyl chain optionally containing 1 or more eg. 2 N, S, O or P chain atoms which can be functionalised, or O, each Y, which may be the same or different, is O or S and each R¹, which may be the same or different, is as defined above,

10




where m is 0 or 1, Q represents (CR¹R⁶)_xZ where x is 0, 1 or 2, R¹ is as defined above and R⁶ is as defined for R¹ and Z is P=O(OH)₂, B(OH)₂ or SO₃H, or a salt thereof, or

15



where each R¹, which are the same or different, is as defined above; R¹¹ represents OH or R¹⁰ NH where R¹⁰ is HO, R¹CO or HOOC(X)_x wherein R¹ is as defined above, x is 0 or 1 and X is R¹R¹C wherein each R¹, which are the same or different, is as defined above; or R¹⁰ is an amino acid residue H₂N (R¹R¹C) CO- wherein each R¹, which are the same or different, is as defined above; n is 1 or 2 and R¹² is H or straight or branched C₁ to C₆ alkyl; or a salt thereof,

25



wherein each of Z¹ and Z² is independently hydrogen, SH or an electron withdrawing group, and R¹² is as defined above, or a salt thereof for use in the treatment of a

30

condition associated with increased or decreased HIF levels or activity, or a condition in which an increase or decrease in HIF levels or activity may be beneficial.

2. A compound according to claim 1 which is of formula (A) where R¹ is aryloxymethyl, R² is hydrogen and R³ is hydrogen.

5 3. A compound according to claim 2 wherein R¹ is phenyloxymethyl, benzyloxymethyl or p-hydroxybenzylmethyl.

4. A compound according to claim 1 of formula (B) where X is a single bond or methyl, R² is H or phenyl alkyl and R⁴ is H or methyl.

10 5. A compound according to claim 1 of formula (C) wherein X is O and R¹ is H or methyl.

6. A compound according to claim 1 of formula (D) wherein x is 1, and R¹ and R⁶ are both hydrogen.

7. A compound according to claim 1 of formula (E) wherein R¹ is H and R¹² is H or ethyl.

15 8. A compound according to claim 7 wherein R¹¹ represents R¹⁰ NH and R¹⁰ is an acyl group of a natural amino acid, acetyl or benzoyl.

9. A compound according to claim 7 wherein R¹¹ is HOOC(X)x.

10. Use of a compound as defined in claim 1 in the manufacture of a medicament for the treatment of a condition associated with increased or decreased HIF levels or activity, or a condition in which an increase or decrease in HIF levels or activity may be beneficial.

11. Use according to claim 10 wherein the compound is as defined in any one of claims 2 to 9.

12. A pharmaceutical composition which comprises a compound as defined in claim 1 together with a pharmaceutically acceptable excipient vehicle or carrier.

13. A composition according to claim 12 wherein the compound is as defined in any one of claims 2 to 9.

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5



SEQUENCE LISTING

<110> ISIS INNOVATION LIMITED
 <120> HIF HYDROXYLASE INHIBITORS
 5 <130> N.85209A PEJ

 <160> 6
 <170> PatentIn version 3.1

 10 <210> 1
 <211> 2110
 <212> DNA
 <213> Homo sapiens
 <220>
 15 <221> CDS
 <222> (297)..(1517)
 <223>

 <400> 1
 20 gctttccct gctgctgt ctctagtttc tctcacatcc cttttttttt tcctttctct 60
 agccaccctg aagggtccct tccaagccc ttagggaccg cagaggactt ggggaccagc 120
 aagcaacccc cagggcacga gaagagctct tgctgtctgc cctgcctcac cctgcccac 180
 accaggcccg gtggcccca gctgcatcaa gtggaggcgg aggaggaggc ggaggagggt 240
 ggcaccatgg gcccgggcgg tgccctccat gcccggggga tgaagacact gctgcc atg 299
 25 Met
 1
 gac agc ccg tgc cag ccg cag ccc cta agt cag gct ctc cct cag tta 347
 Asp Ser Pro Cys Gln Pro Gln Pro Leu Ser Gln Ala Leu Pro Gln Leu
 5 10 15
 30 cca ggg tct tcg tca gag ccc ttg gag cct gag cct ggc cgg gcc agg 395
 Pro Gly Ser Ser Ser Glu Pro Leu Glu Pro Glu Pro Gly Arg Ala Arg
 20 25 30
 atg gga gtg gag agt tac ctg ccc tgt ccc ctg ctc ccc tcc tac cac 443
 Met Gly Val Glu Ser Tyr Leu Pro Cys Pro Leu Leu Pro Ser Tyr His
 35 35 40 45
 35 tgt cca gga gtg cct agt gag gcc tcg gca ggg agt ggg acc ccc aga 491
 Cys Pro Gly Val Pro Ser Glu Ala Ser Ala Gly Ser Gly Thr Pro Arg
 50 55 60 65
 40 gcc aca gcc acc tct acc act gcc agc cct ctt cgg gac ggt ttt ggc 539
 Ala Thr Ala Thr Ser Thr Thr Ala Ser Pro Leu Arg Asp Gly Phe Gly
 70 75 80
 ggg cag gat ggt ggt gag ctg cgg ccg ctg cag agt gaa ggc gct gca 587
 Gly Gln Asp Gly Gly Glu Leu Arg Pro Leu Gln Ser Glu Gly Ala Ala
 85 90 95
 45 gcg ctg gtc acc aag ggg tgc cag cga ttg gca gcc cag ggc gca cgg 635
 Ala Leu Val Thr Lys Gly Cys Gln Arg Leu Ala Ala Gln Gly Ala Arg
 100 105 110
 cct gag gcc ccc aaa cgg aaa tgg gcc gag gat ggt ggg gat gcc cct 683
 Pro Glu Ala Pro Lys Arg Lys Trp Ala Glu Asp Gly Gly Asp Ala Pro

	115	120	125	
	tca ccc agc aaa cgg ccc tgg gcc agg caa gag aac cag gag gca gag			731
	Ser Pro Ser Lys Arg Pro Trp Ala Arg Gln Glu Asn Gln Glu Ala Glu			
	130	135	140	145
5	cgg gag ggt ggc atg agc tgc agc tgc agc agt ggc agt ggt gag gcc			779
	Arg Glu Gly Gly Met Ser Cys Ser Cys Ser Ser Gly Ser Gly Glu Ala			
	150	155	160	
	agt gct ggg ctg atg gag gag gcg ctg ccc tct gcg ccc gag cgc ctg			827
	Ser Ala Gly Leu Met Glu Glu Ala Leu Pro Ser Ala Pro Glu Arg Leu			
10	165	170	175	
	gcc ctg gac tat atc gtg ccc tgc atg cgg tac tac ggc atc tgc gtc			875
	Ala Leu Asp Tyr Ile Val Pro Cys Met Arg Tyr Tyr Gly Ile Cys Val			
	180	185	190	
15	aag gac agc ttc ctg ggg gca gca ctg ggc ggt cgc gtg ctg gcc gag			923
	Lys Asp Ser Phe Leu Gly Ala Ala Leu Gly Gly Arg Val Leu Ala Glu			
	195	200	205	
	gtg gag gcc ctc aaa cgg ggt ggg cgc ctg cga gac ggg cag cta gtg			971
	Val Glu Ala Leu Lys Arg Gly Gly Arg Leu Arg Asp Gly Gln Leu Val			
	210	215	220	225
20	agc cag agg gcg atc ccg ccg cgc agc atc cgt ggg gac cag att gcc			1019
	Ser Gln Arg Ala Ile Pro Pro Arg Ser Ile Arg Gly Asp Gln Ile Ala			
	230	235	240	
	tgg gtg gaa ggc cat gaa cca ggc tgt cga agc att ggt gcc ctc atg			1067
	Trp Val Glu Gly His Glu Pro Gly Cys Arg Ser Ile Gly Ala Leu Met			
25	245	250	255	
	gcc cat gtg gac gcc gtc atc cgc cac tgc gca ggg cgg ctg ggc agc			1115
	Ala His Val Asp Ala Val Ile Arg His Cys Ala Gly Arg Leu Gly Ser			
	260	265	270	
	tat gtc atc aac ggg cgc acc aag gcc atg gtg gcg tgt tac cca ggc			1163
30	Tyr Val Ile Asn Gly Arg Thr Lys Ala Met Val Ala Cys Tyr Pro Gly			
	275	280	285	
	aac ggg ctc ggg tac gta agg cac gtt gac aat ccc cac ggc gat ggg			1211
	Asn Gly Leu Gly Tyr Val Arg His Val Asp Asn Pro His Gly Asp Gly			
	290	295	300	305
35	cgc tgc atc acc tgt atc tat tac ctg aat cag aac tgg gac gtt aag			1259
	Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Gln Asn Trp Asp Val Lys			
	310	315	320	
	gtg cat ggc ggc ctg ctg cag atc ttc cct gag ggc cgg ccc gtg gta			1307
	Val His Gly Gly Leu Leu Gln Ile Phe Pro Glu Gly Arg Pro Val Val			
40	325	330	335	
	gcc aac atc gag cca ctc ttt gac cgg ttg ctc att ttc tgg tct gac			1355
	Ala Asn Ile Glu Pro Leu Phe Asp Arg Leu Leu Ile Phe Trp Ser Asp			
	340	345	350	
	cgg cgg aac ccc cac gag gtg aag cca gcc tat gcc acc agg tac gcc			1403
45	Arg Arg Asn Pro His Glu Val Lys Pro Ala Tyr Ala Thr Arg Tyr Ala			
	355	360	365	
	atc act gtc tgg tat ttt gat gcc aag gag cgg gca gca gcc aaa gac			1451
	Ile Thr Val Trp Tyr Phe Asp Ala Lys Glu Arg Ala Ala Ala Lys Asp			
	370	375	380	385

	aag tat cag cta gca tca gga cag aaa ggt gtc caa gta cct gta tca	1499
	Lys Tyr Gln Leu Ala Ser Gly Gln Lys Gly Val Gln Val Pro Val Ser	
	390 395 400	
5	cag ccg cct acg ccc acc tagtgccag tcccagagcc gcatggcaga	1547
	Gln Pro Pro Thr Pro Thr	
	405	
	cagcttaaat gacttcagga gagccctggg cctgtgctgg ctgctccttc cctgccaccg	1607
	ctgctgcttc tgactttgcc tctgtcctgc ctgggtgga gggctctgtc tgttgtgag	1667
	gaccaaggag gagaagagac ctttctgcc ccatcatggg ggctggggtt gtcacctgga	1727
10	cagggggcag ccgtggaggc caccgttacc aactgaagct gggggcctgg gtcctaccct	1787
	gtctgtgcat gacccatta ggtatggaga gctgggagga ggcattgtca cttcccacca	1847
	ggatgcagga cttgggggtt aggtgagtca tggcctcttg ctggcaatgg ggtgggagga	1907
	gtaccccaa gtctctcac tcctccagcc tggaatgtga agtgactccc caacccttt	1967
	ggccatggca ggcacctttt ggactgggct gccactgctt gggcagagta aaaggtgcca	2027
15	ggaggagcat ggggtgtgaa gtcctgtcag ccaagaaata aaagtttacc tcagagctgc	2087
	aaaaaaaa aaaaaaaaa aaa	2110
	<210> 2	
	<211> 407	
20	<212> PRT	
	<213> Homo sapiens	
	<400> 2	
	Met Asp Ser Pro Cys Gln Pro Gln Pro Leu Ser Gln Ala Leu Pro Gln	
25	1 5 10 15	
	Leu Pro Gly Ser Ser Ser Glu Pro Leu Glu Pro Glu Pro Gly Arg Ala	
	20 25 30	
	Arg Met Gly Val Glu Ser Tyr Leu Pro Cys Pro Leu Leu Pro Ser Tyr	
	35 40 45	
30	His Cys Pro Gly Val Pro Ser Glu Ala Ser Ala Gly Ser Gly Thr Pro	
	50 55 60	
	Arg Ala Thr Ala Thr Ser Thr Thr Ala Ser Pro Leu Arg Asp Gly Phe	
	65 70 75 80	
	Gly Gly Gln Asp Gly Gly Glu Leu Arg Pro Leu Gln Ser Glu Gly Ala	
35	85 90 95	
	Ala Ala Leu Val Thr Lys Gly Cys Gln Arg Leu Ala Ala Gln Gly Ala	
	100 105 110	
	Arg Pro Glu Ala Pro Lys Arg Lys Trp Ala Glu Asp Gly Gly Asp Ala	
	115 120 125	
40	Pro Ser Pro Ser Lys Arg Pro Trp Ala Arg Gln Glu Asn Gln Glu Ala	
	130 135 140	
	Glu Arg Glu Gly Gly Met Ser Cys Ser Cys Ser Ser Gly Ser Gly Glu	
	145 150 155 160	
	Ala Ser Ala Gly Leu Met Glu Glu Ala Leu Pro Ser Ala Pro Glu Arg	
45	165 170 175	
	Leu Ala Leu Asp Tyr Ile Val Pro Cys Met Arg Tyr Tyr Gly Ile Cys	
	180 185 190	
	Val Lys Asp Ser Phe Leu Gly Ala Ala Leu Gly Gly Arg Val Leu Ala	
	195 200 205	

Glu Val Glu Ala Leu Lys Arg Gly Gly Arg Leu Arg Asp Gly Gln Leu
 210 215 220
 Val Ser Gln Arg Ala Ile Pro Pro Arg Ser Ile Arg Gly Asp Gln Ile
 225 230 235 240
 5 Ala Trp Val Glu Gly His Glu Pro Gly Cys Arg Ser Ile Gly Ala Leu
 245 250 255
 Met Ala His Val Asp Ala Val Ile Arg His Cys Ala Gly Arg Leu Gly
 260 265 270
 10 Ser Tyr Val Ile Asn Gly Arg Thr Lys Ala Met Val Ala Cys Tyr Pro
 275 280 285
 Gly Asn Gly Leu Gly Tyr Val Arg His Val Asp Asn Pro His Gly Asp
 290 295 300
 Gly Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Gln Asn Trp Asp Val
 305 310 315 320
 15 Lys Val His Gly Gly Leu Leu Gln Ile Phe Pro Glu Gly Arg Pro Val
 325 330 335
 Val Ala Asn Ile Glu Pro Leu Phe Asp Arg Leu Leu Ile Phe Trp Ser
 340 345 350
 20 Asp Arg Arg Asn Pro His Glu Val Lys Pro Ala Tyr Ala Thr Arg Tyr
 355 360 365
 Ala Ile Thr Val Trp Tyr Phe Asp Ala Lys Glu Arg Ala Ala Ala Lys
 370 375 380
 Asp Lys Tyr Gln Leu Ala Ser Gly Gln Lys Gly Val Gln Val Pro Val
 385 390 395 400
 25 Ser Gln Pro Pro Thr Pro Thr
 405

<210> 3
 <211> 5163
 30 <212> DNA
 <213> Homo sapiens
 <220>

<220>
 35 <221> CDS
 <222> (3157)..(4434)
 <223>

<400> 3
 40 ttaggggcag aaaaacattt gtaataatta atggctttga gagacacaag gctttgtttg 60
 cccagagta ttagttaacc cacctagtgc tcctaatacat acaatattaa ggattgggag 120
 ggacattcat tgcctcactc tctatttgtt tcaccttctg taaaattggt agaataatag 180
 taccacttc atagcattgt atgatgatta aattggttaa tatttttaaa atgcttagaa 240
 cacagattgg gcacataaca gcaagcacca catgtgttta taagataaat tcctttgtgt 300
 45 tgccttcctg taaagttaa ataagtaaat aaataaataa atacttgc atgacattttga 360
 agtctctcta taacatctga gtaagtggcg gctgcgacaa tgctactgga gttccagaat 420
 cgtgttggtg acaagattgt tcaccagcat atggtgtggt gaaaactcac taatttgga 480
 ttagttcaga ttattaagcc tgaatagggtg aaaaatcctga aatcaaggat ctttggaact 540
 atttgaaatc agtattttat attttcctgt tgtattcatt aaagtgttgc aagtgttcta 600

	ttt gat g gat	taagt at att	tagga ta tac	atgtt ca att	tgt gat ttt	tata ct ta at	660
	tgga a ca aga	aag ct a ta aa	agg ttt t gat	atgg a cat ct	att ct ttt aa	gtaa a ct t ca	720
	atgaaa at at	atg ag tag ag	cat at ag ag a	tgta a at a at	ttgt gg a ca c	acc a ca g act	780
	gaa at ag ca a	at t t t a a a ga	aatt gtt gga	aga at ca agt	gttt gt gg aa	tgag t cct cc	840
5	tagt aa gtt	cct gct ct t g	tga at a att a	agc ct cat gt	ata att act a	tag ca a a ag g	900
	aag cta aga	agt att ag ac	tct act t g ta	ttta a att ac	at ttt ac at a	at ttt at g t g	960
	atgaaa at g	tt t t a a at gc	ttatt t t cgt	aag cc at g ag	at ag ct cct t	tat att ttt aa	1020
	gaatt t ct ga	at t a att t g c	ttgg att tta	ttag t g ca aa	tg gc ag ag ct	ag ca att c ct	1080
	tttt ct g t g	tccc att tcca	tcct att cat	ccct ctt tta	ggaa act ct g	aact ct gg at	1140
10	tg t cct t gtt	tac at ac ct g	cct cct g cat	tgg act at gt	gt ct ct g ag t	gtag tat g ac	1200
	taatt cat ttt	gttt gt ca ag	gact ct ca at	gcatt t gtt g	aac ag cct aa	ttag ta at gt	1260
	ctg ca a ca at	gac att ttt ac	tgt att tta at	aa ag ct ct gg	gaa ag tag ga	tac ac at a ag	1320
	ac ag gt ct ag	gt ct aa att c	ttt ac ag aa a	ctt gg att ttt	tag ttc ggt t	tga a att t g a	1380
	ag at gt g ag t	at att ttt at ct	cag ttt tcca	aag gac a ag c	ta att gga at	tat cat cct c	1440
15	ttt cact tga	ttgg at cccc	aga at gcc at	ttac g cat gc	agc ag g att t	tata ac ag tt	1500
	ttaa att ct g	tat att t g at	ga ag ag gtt t	tat att ttt t g	gatt ca ag cc	tct ttt tta aa	1560
	ctt ct a ca at	at ggt ttt aca	ata att cct t	at at cct g ct	ttt gaa at ac	at att a ca ac	1620
	tttt a ag tt	tg ga ag g ct a	tatt t ca ag g	act ga ag tta	cag ta act c	aag t ga ta ca	1680
	ca ag ct ag c	acccc act ttt	ccac at ag t g	ttc ga ta a ag	att ga ta a ac	tc ga a at ca c	1740
20	ag ac ttt tta	att ct tta aga	caa at ag c ag	caga a a ga aa	cat ct ttt ggc	ttatt t ct g g	1800
	ta ag gtt ttt	at gct ct g ta	aa a ca a ga a	ttgt att cat	ccg cg cag ca	cag att ct at	1860
	taaaa at a aa	tgt g ag ag t c	gtt a at g tag	tact g ct cat	ttac cat ca a	aatt cact t t	1920
	tc ag ga at aa	tccc at c ag t	ttaa att gga	tatt gga at g	agc att g att	ac att tta act	1980
	tggt ag ccca	aaatt t tct c	at ggg gtt ttt	ga act cg gc g	ggatt t ca aa	ggt ttt taa aa	2040
25	atg ag ttt ttt	gatt ttt ttt t	aaa acc ct ca	aatt t cat ta	cct tta a act	aggt cg aa ac	2100
	gggg cg ca ag	ag att gg att	aac acc at ag	ta act ttt at	ttt gtt ct ta	acc att t c ag	2160
	ggct tct tga	aat ag ag gct	gt at ggt g ta	at gga a a a a	cag cct t gga	at ct ggg ag c	2220
	ctg att cct g	gatt cag tcc	cag ttt t g c g	tg ac ct t g g g	ca ag tta ct t	tact tct ct g	2280
	aatt tcc gtt	tcct cct ct g	caaa at g ag g	atc g ca at ag	ccac ct t g ca	ac ct t g act g	2340
30	gag cg ag cct	cgc ac a cccc	gcg cc ggc ct	gg ag ga ag ag	cag cc at g at	tac g cc g cct	2400
	tcg ct cc g et	accc g ct t g c	gg ct g g c g cc	ct cct cc ag c	agg t g ta g g c	gct g cc g c g c	2460
	tgcccc ac g c	cttt cc g cc g	ctc g c g g g cc	tg c g c ct g g	cgt cccc g ag	gagg cc g ct g	2520
	cgg gct g ag g	tag c g ca cc g	gcct ct c g g c	gt ccc ag tcc	ggt ccc g g g c	gg ag g g a a g	2580
	cgg gc ga ccc	ac ct cc g ag g	caga ag cc ga	gg cc c g g c c c	cg cc g ag t g c	gg ag g ag c g c	2640
35	agg ca g cccc	cg ccc ct c g g	cc ct c c c c c c	gg cc ct c c c g	gcc ct c c c t c	cg ccc c c t c c	2700
	gcc ct c g c g c	gcc g c c c g c c	cgg g t c g c c g	cgg g g c c g t g	gt g ta c g t g c	ag ag c g c g ca	2760
	gag cg ag t g g	cg ccc g ta t g	cc ct g c g t c	ct cca ca g cc	tgg g c c g g g c	cg ccc g g g ac	2820
	gct g ag g c g g	cg g c g g c g c	cg ag g g g g c c	ggt ct t g c g c	tccc c ag g c c	cg c g c g c t g	2880
	ag ccc ag g tt	gcc att c g c c	gc ac ag g c c c	tatt ct ct ca	gcc ct c g g c g	gc gat g ag g c	2940
40	gct g ag g c g g	ct g c c g g c g c	tg c g c c g g ag	ct tag g act c	gga ag c g g c c	gg g c c g ag g g	3000
	cgt g g g g t g c	cg g c ct c c ct	gag g c g ag g g	tag c g g g t g c	at g g c g c ag t	aac g g c c c ct	3060
	at ct ct ct c c	cc g ct c c c ca	gc ct c g g g c g	agg cc g t c c g	gcc g ct ac c c	ct c ct g ct c g	3120
	gcc g c c g c ag	tc g c g t c g c	cg cc g c c g c c	gcc g c c at g	gcc aat gac agc ggc		3174
				Met Ala Asn Asp Ser Gly			
45				1	5		
	ggg ccc ggc	ggg ccg agc	ccg agc gag	cga gac	cgg cag tac	tgc gag	3222
	Gly Pro Gly	Gly Pro Ser	Pro Ser Glu	Arg Asp	Arg Gln Tyr	Cys Glu	
		10	15		20		
	ctg tgc ggg	aag atg gag	aac ctg ctg	cgc tgc	agc cgc tgc	cgc agc	3270

	Leu Cys Gly Lys Met Glu Asn Leu Leu Arg Cys Ser Arg Cys Arg Ser	
	25 30 35	
5	tcc ttc tac tgc tgc aag gag cac cag cgt cag gac tgg aag aag cac Ser Phe Tyr Cys Cys Lys Glu His Gln Arg Gln Asp Trp Lys Lys His	3318
	40 45 50	
	aag ctc gtg tgc cag ggc agc gag ggc gcc ctc ggc cac gga gtg ggc Lys Leu Val Cys Gln Gly Ser Glu Gly Ala Leu Gly His Gly Val Gly	3366
	55 60 65 70	
10	cca cac cag cat tcc ggc ccc gcg ccg ccg gct gca gtg ccg ccg ccc Pro His Gln His Ser Gly Pro Ala Pro Pro Ala Ala Val Pro Pro Pro	3414
	75 80 85	
	agg gcc ggg gcc cgg gag ccc agg aag gca gcg gcg cgc ccg gac aac Arg Ala Gly Ala Arg Glu Pro Arg Lys Ala Ala Ala Arg Arg Asp Asn	3462
	90 95 100	
15	gcc tcc ggg gac gcg gcc aag gga aaa gta aag gcc aag ccc ccg gcc Ala Ser Gly Asp Ala Ala Lys Gly Lys Val Lys Ala Lys Pro Pro Ala	3510
	105 110 115	
	gac cca gcg gcg gcc gcg tcg ccg tgt cgt gcg gcc gcc ggc ggc cag Asp Pro Ala Ala Ala Ala Ser Pro Cys Arg Ala Ala Ala Gly Gly Gln	3558
20	120 125 130	
	ggc tcg gcg gtg gct gcc gaa gcc gag ccc ggc aag gag gag ccg ccg Gly Ser Ala Val Ala Ala Glu Ala Glu Pro Gly Lys Glu Glu Pro Pro	3606
	135 140 145 150	
25	gcc cgc tca tcg ctg ttc cag gag aag gcg aac ctg tac ccc cca agc Ala Arg Ser Ser Leu Phe Gln Glu Lys Ala Asn Leu Tyr Pro Pro Ser	3654
	155 160 165	
	aac acg ccc ggg gat gcg ctg agc ccc ggc ggc ggc ctg ccg ccc aac Asn Thr Pro Gly Asp Ala Leu Ser Pro Gly Gly Gly Leu Arg Pro Asn	3702
	170 175 180	
30	ggg cag acg aag ccc ctg ccg gcg ctg aag ctg gcg ctc gag tac atc Gly Gln Thr Lys Pro Leu Pro Ala Leu Lys Leu Ala Leu Glu Tyr Ile	3750
	185 190 195	
	gtg ccg tgc atg aac aag cac ggc atc tgt gtg gtg gac gac ttc ctc Val Pro Cys Met Asn Lys His Gly Ile Cys Val Val Asp Asp Phe Leu	3798
35	200 205 210	
	ggc aag gag acc gga cag cag atc ggc gac gag gtg cgc gcc ctg cac Gly Lys Glu Thr Gly Gln Gln Ile Gly Asp Glu Val Arg Ala Leu His	3846
	215 220 225 230	
40	gac acc ggg aag ttc acg gac ggg cag ctg gtc agc cag aag agt gac Asp Thr Gly Lys Phe Thr Asp Gly Gln Leu Val Ser Gln Lys Ser Asp	3894
	235 240 245	
	tcg tcc aag gac atc cga ggc gat aag atc acc tgg atc gag ggc aag Ser Ser Lys Asp Ile Arg Gly Asp Lys Ile Thr Trp Ile Glu Gly Lys	3942
	250 255 260	
45	gag ccc ggc tgc gaa acc att ggg ctg ctc atg agc agc atg gac gac Glu Pro Gly Cys Glu Thr Ile Gly Leu Leu Met Ser Ser Met Asp Asp	3990
	265 270 275	
	ctg ata cgc cac tgt aac ggg aag ctg ggc agc tac aaa atc aat ggc Leu Ile Arg His Cys Asn Gly Lys Leu Gly Ser Tyr Lys Ile Asn Gly	4038

	280	285	290	
	cgg acg aaa gcc atg gtt gct tgt tat ccg ggc aat gga acg ggt tat			4086
	Arg Thr Lys Ala Met Val Ala Cys Tyr Pro Gly Asn Gly Thr Gly Tyr			
	295	300	305	310
5	gta cgt cat gtt gat aat cca aat gga gat gga aga tgt gtg aca tgt			4134
	Val Arg His Val Asp Asn Pro Asn Gly Asp Gly Arg Cys Val Thr Cys			
	315	320	325	
	ata tat tat ctt aat aaa gac tgg gat gcc aag gta agt gga ggt ata			4182
	Ile Tyr Tyr Leu Asn Lys Asp Trp Asp Ala Lys Val Ser Gly Gly Ile			
10	330	335	340	
	ctt cga att ttt cca gaa ggc aaa gcc cag ttt gct gac att gaa ccc			4230
	Leu Arg Ile Phe Pro Glu Gly Lys Ala Gln Phe Ala Asp Ile Glu Pro			
	345	350	355	
	aaa ttt gat aga ctg ctg ttt ttc tgg tct gac cgt cgc aac cct cat			4278
15	Lys Phe Asp Arg Leu Leu Phe Phe Trp Ser Asp Arg Arg Asn Pro His			
	360	365	370	
	gaa gta caa cca gca tat gct aca agg tac gca ata act gtt tgg tat			4326
	Glu Val Gln Pro Ala Tyr Ala Thr Arg Tyr Ala Ile Thr Val Trp Tyr			
	375	380	385	390
20	ttt gat gca gat gag aga gca cga gct aaa gta aaa tat cta aca ggt			4374
	Phe Asp Ala Asp Glu Arg Ala Arg Ala Lys Val Lys Tyr Leu Thr Gly			
	395	400	405	
	gaa aaa ggt gtg agg gtt gaa ctc aat aaa cct tca gat tcg gtc ggt			4422
	Glu Lys Gly Val Arg Val Glu Leu Asn Lys Pro Ser Asp Ser Val Gly			
25	410	415	420	
	aaa gac gtc ttc tagagccttt gatccagcaa taccccactt cacctacaat			4474
	Lys Asp Val Phe			
	425			
	attgttaact atttgttaac ttgtgaatac gaataaatgg gataaagaaa aatagacaac			4534
30	cagttcgcac ttttaataagg aaacagaaac aactttttgt gttgcatcaa acagaagatt			4594
	ttgactgctg tgacttttga ctgcatgac aacttcaaat ctgtgattgc ttacaggagg			4654
	aagataagct actaattgaa aatgggtttt acatctggat atgaaataag tgccctgtgt			4714
	agaatttttt tcattcttat attttgccag atctgttatc tagctgagtt catttcatct			4774
	ctcccttttt tatatcaagt ttgaatttgg gataattttt ctatattagg tacaatttat			4834
35	ctaaactgaa ttgagaaaaa attacagtat tattcctcaa aataacatca atctattttt			4894
	gtaaacctgt tcatactatt aaattttgcc ctaaaagacc tcttaataat gattgttgcc			4954
	agtgactgat gattaatttt attttactta aaataagaaa aggagcactt taattacaac			5014
	tgaaaaatca gattgttttg cagtccttcc ttacactaat ttgaactctt aaagattgct			5074
	gctttttttt tgacattgtc aataacgaaa cctaattgta aaacagtcac catttactac			5134
40	caataacttt tagttaatgt tttaacaagg			5163
	<210> 4			
	<211> 426			
	<212> PRT			
45	<213> Homo sapiens			
	<400> 4			
	Met Ala Asn Asp Ser Gly Gly Pro Gly Gly Pro Ser Pro Ser Glu Arg			
	1 5 10 15			

	Asp	Arg	Gln	Tyr	Cys	Glu	Leu	Cys	Gly	Lys	Met	Glu	Asn	Leu	Leu	Arg
			20						25				30			
	Cys	Ser	Arg	Cys	Arg	Ser	Ser	Phe	Tyr	Cys	Cys	Lys	Glu	His	Gln	Arg
		35						40					45			
5	Gln	Asp	Trp	Lys	Lys	His	Lys	Leu	Val	Cys	Gln	Gly	Ser	Glu	Gly	Ala
	50						55					60				
	Leu	Gly	His	Gly	Val	Gly	Pro	His	Gln	His	Ser	Gly	Pro	Ala	Pro	Pro
	65					70					75				80	
	Ala	Ala	Val	Pro	Pro	Pro	Arg	Ala	Gly	Ala	Arg	Glu	Pro	Arg	Lys	Ala
10					85					90					95	
	Ala	Ala	Arg	Arg	Asp	Asn	Ala	Ser	Gly	Asp	Ala	Ala	Lys	Gly	Lys	Val
			100						105				110			
	Lys	Ala	Lys	Pro	Pro	Ala	Asp	Pro	Ala	Ala	Ala	Ala	Ser	Pro	Cys	Arg
			115					120					125			
15	Ala	Ala	Ala	Gly	Gly	Gln	Gly	Ser	Ala	Val	Ala	Ala	Glu	Ala	Glu	Pro
	130						135					140				
	Gly	Lys	Glu	Glu	Pro	Pro	Ala	Arg	Ser	Ser	Leu	Phe	Gln	Glu	Lys	Ala
	145					150					155				160	
	Asn	Leu	Tyr	Pro	Pro	Ser	Asn	Thr	Pro	Gly	Asp	Ala	Leu	Ser	Pro	Gly
20					165					170					175	
	Gly	Gly	Leu	Arg	Pro	Asn	Gly	Gln	Thr	Lys	Pro	Leu	Pro	Ala	Leu	Lys
			180						185					190		
	Leu	Ala	Leu	Glu	Tyr	Ile	Val	Pro	Cys	Met	Asn	Lys	His	Gly	Ile	Cys
		195						200					205			
25	Val	Val	Asp	Asp	Phe	Leu	Gly	Lys	Glu	Thr	Gly	Gln	Gln	Ile	Gly	Asp
	210						215					220				
	Glu	Val	Arg	Ala	Leu	His	Asp	Thr	Gly	Lys	Phe	Thr	Asp	Gly	Gln	Leu
	225					230					235				240	
	Val	Ser	Gln	Lys	Ser	Asp	Ser	Ser	Lys	Asp	Ile	Arg	Gly	Asp	Lys	Ile
30					245					250					255	
	Thr	Trp	Ile	Glu	Gly	Lys	Glu	Pro	Gly	Cys	Glu	Thr	Ile	Gly	Leu	Leu
		260						265					270			
	Met	Ser	Ser	Met	Asp	Asp	Leu	Ile	Arg	His	Cys	Asn	Gly	Lys	Leu	Gly
		275						280					285			
35	Ser	Tyr	Lys	Ile	Asn	Gly	Arg	Thr	Lys	Ala	Met	Val	Ala	Cys	Tyr	Pro
	290					295						300				
	Gly	Asn	Gly	Thr	Gly	Tyr	Val	Arg	His	Val	Asp	Asn	Pro	Asn	Gly	Asp
	305					310					315				320	
	Gly	Arg	Cys	Val	Thr	Cys	Ile	Tyr	Tyr	Leu	Asn	Lys	Asp	Trp	Asp	Ala
40					325					330					335	
	Lys	Val	Ser	Gly	Gly	Ile	Leu	Arg	Ile	Phe	Pro	Glu	Gly	Lys	Ala	Gln
			340						345					350		
	Phe	Ala	Asp	Ile	Glu	Pro	Lys	Phe	Asp	Arg	Leu	Leu	Phe	Phe	Trp	Ser
		355						360					365			
45	Asp	Arg	Arg	Asn	Pro	His	Glu	Val	Gln	Pro	Ala	Tyr	Ala	Thr	Arg	Tyr
	370						375					380				
	Ala	Ile	Thr	Val	Trp	Tyr	Phe	Asp	Ala	Asp	Glu	Arg	Ala	Arg	Ala	Lys
	385					390					395				400	
	Val	Lys	Tyr	Leu	Thr	Gly	Glu	Lys	Gly	Val	Arg	Val	Glu	Leu	Asn	Lys

	aag aat tgg gat gcc aag cta cat ggt ggg atc ctg cgg ata ttt cca	833
	Lys Asn Trp Asp Ala Lys Leu His Gly Gly Ile Leu Arg Ile Phe Pro	
	155 160 165	
5	gag ggg aaa tca ttc ata gca gat gtg gag ccc att ttt gac aga ctc	881
	Glu Gly Lys Ser Phe Ile Ala Asp Val Glu Pro Ile Phe Asp Arg Leu	
	170 175 180 185	
	ctg ttc ttc tgg tca gat cgt agg aac cca cac gaa gtg cag ccc tct	929
	Leu Phe Phe Trp Ser Asp Arg Arg Asn Pro His Glu Val Gln Pro Ser	
	190 195 200	
10	tac gca acc aga tat gct atg act gtc tgg tac ttt gat gct gaa gaa	977
	Tyr Ala Thr Arg Tyr Ala Met Thr Val Trp Tyr Phe Asp Ala Glu Glu	
	205 210 215	
	agg gca gaa gcc aaa aag aaa ttc agg aat tta act agg aaa act gaa	1025
	Arg Ala Glu Ala Lys Lys Lys Phe Arg Asn Leu Thr Arg Lys Thr Glu	
15	220 225 230	
	tct gcc ctc act gaa gac tgaccgtgct ctgaaatctg ctggccttgt	1073
	Ser Ala Leu Thr Glu Asp	
	235	
20	tcatttttagt aacggttctt gaatttctctt aaatttcttg agatccaaag atggcctctt	1133
	cagtgacaac aatctccctg ctacttcttg catccttcac atccctgtct tgtgtgtggt	1193
	acttcatgtt ttcttgccaa gactgtgttg atcttcagat actctctttg ccagatgaag	1253
	ttatttgcta actccagaaa ttcttgacaga catcctactc ggccagcggg ttacctgata	1313
	gattcggtaa tactatcaag agaagagcct aggagcacag cgagggaatg aaccttactt	1373
	gcactttatg tatacttcct gatttgaaag gaggaggttt gaaaagaaaa aaatggaggt	1433
25	ggtagatgcc acagagaggc atcacggaag ccttaacagc aggaacaga gaaattgtg	1493
	tcatctgaac aatttccaga tgttcttaat ccagggtgtg tggggtttct ggagaattat	1553
	cacaacctaa tgacattaat acctctagaa agggctgctg tcatagttaa caatttataa	1613
	gtgtcccatg gggcagacac tccttttttc ccagtctgc aacctggatt ttctgcctca	1673
	gctccatttt gctgaaaata atgactttct gaataaagat ggcaacacaa ttttttctcc	1733
30	attttcagtt cttacctggg aacctaatc cccagaagct aaaaaactag acattagtgtg	1793
	ttttggttg tttgttgaa tggaatttaa atttaaatga aaggaaaaat atatccctgg	1853
	tagttttgtg ttaacctag ataactgtgg aaagagctag gtctactgat atacaataaa	1913
	catgtgtgca tcttgaacaa tttagagagg gaggtggagt tggaatgtg ggtgttctctg	1973
	ttttttttt ttttttttt ttttttttagt ttctctttt aatgagctca ccctttaaca	2033
35	caaaaaaagc agggatgatgt attttaaaaa aggaagtgga aataaaaaaa tctcaaagct	2093
	atttgagttc tcgtctgtcc cttagcagct ttcttcagct cacttggctc tctagatcca	2153
	ctgtggttgg cagtatgacc agaatcatgg aacttgctag aactgtggaa gcttctactc	2213
	ctgcagtaag cacagatcgc actgcctcaa taacttggtg ttgagcacgt attttgcaaa	2273
	agctactttt cctagttttc agtattactt tcatgtttta aaaatccctt taatttcttg	2333
40	cttgaaaatc ccatgaacat taaagagcca gaaatatttt cctttgttat gtacggatat	2393
	atatatatat atagtcttcc aagatagaag ttacttttt cctcttcttg ttttgaaaaa	2453
	tttccagata agacatgtca ccattaatc tcaacgactg ctctatatttg ttgtacggta	2513
	atagttatca ctttctaaat tactatgtaa ttactcact tattatgttt attgtcttgt	2573
	atcctttctc tggagtgtaa gcacaatgaa gacaggaatt ttgtatatat ttaaccaatg	2633
45	caacatactc tcagcaccta aaatagtgcc gggaacatag taagggtcga gtaaatactt	2693
	gttgaataaaa ctacgtctcc tacattagca ttctaaaaaa aaaaaaaaaa aaaaaaaaaa	2753
	aaaaaaaaa aaaaaaag	2770

<211> 239
 <212> PRT
 <213> Homo sapiens
 <400> 6

5
 Met Pro Leu Gly His Ile Met Arg Leu Asp Leu Glu Lys Ile Ala Leu
 1 5 10 15
 Glu Tyr Ile Val Pro Cys Leu His Glu Val Gly Phe Cys Tyr Leu Asp
 20 25 30
 10 Asn Phe Leu Gly Glu Val Val Gly Asp Cys Val Leu Glu Arg Val Lys
 35 40 45
 Gln Leu His Cys Thr Gly Ala Leu Arg Asp Gly Gln Leu Ala Gly Pro
 50 55 60
 Arg Ala Gly Val Ser Lys Arg His Leu Arg Gly Asp Gln Ile Thr Trp
 15 65 70 75 80
 Ile Gly Gly Asn Glu Glu Gly Cys Glu Ala Ile Ser Phe Leu Leu Ser
 85 90 95
 Leu Ile Asp Arg Leu Val Leu Tyr Cys Gly Ser Arg Leu Gly Lys Tyr
 100 105 110
 20 Tyr Val Lys Glu Arg Ser Lys Ala Met Val Ala Cys Tyr Pro Gly Asn
 115 120 125
 Gly Thr Gly Tyr Val Arg His Val Asp Asn Pro Asn Gly Asp Gly Arg
 130 135 140
 Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Lys Asn Trp Asp Ala Lys Leu
 145 150 155 160
 25 His Gly Gly Ile Leu Arg Ile Phe Pro Glu Gly Lys Ser Phe Ile Ala
 165 170 175
 Asp Val Glu Pro Ile Phe Asp Arg Leu Leu Phe Phe Trp Ser Asp Arg
 180 185 190
 30 Arg Asn Pro His Glu Val Gln Pro Ser Tyr Ala Thr Arg Tyr Ala Met
 195 200 205
 Thr Val Trp Tyr Phe Asp Ala Glu Glu Arg Ala Glu Ala Lys Lys Lys
 210 215 220
 Phe Arg Asn Leu Thr Arg Lys Thr Glu Ser Ala Leu Thr Glu Asp
 35 225 230 235

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.